



Results from FOPI on nuclear collective flow in heavy ion collisions at SIS energies

N. Bastid

► To cite this version:

N. Bastid. Results from FOPI on nuclear collective flow in heavy ion collisions at SIS energies. International Nuclear Physics Conferences (INPC 2004) 1, Jun 2004, Goteborg, Sweden. pp.1-12. in2p3-00022230

HAL Id: in2p3-00022230

<https://hal.in2p3.fr/in2p3-00022230>

Submitted on 9 Sep 2004

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.



L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Results from FOPI on Nuclear Collective Flow



in Heavy Ion Collisions at SIS energies



1 Motivations

2 FOPI detector overview

3 Experimental systematics

- Directed flow
- Elliptic flow

4 Data *versus* IQMD

- Sensitivity to σ_{mn} ?
- Sensitivity to EoS?

5 Anisotropic flow from Lee-Yang Zeroes

6 Conclusion




Ca + Ca, Ni + Ni, Ru + Ru, Xe + CsI, Au + Au
90A MeV - 2A GeV

Motivations & Observables

Probing hot & dense hadronic matter

↪ Nuclear Equation of State

- Collision dynamics
- In-medium effects: σ_{nn} , MDI









Global flow : $\mathbf{p}_x^{dir} = \sum \text{sign}(y_{cm}) Z u_x / \sum Z, \quad u_x = \beta_x \gamma$


Flow angle: θ_F , Aspect ratios: λ_{31} & λ_{21}

Differential flow : $\frac{dN}{d\varphi'} \sim 1 + 2\mathbf{v}_1 \cos(\varphi') + 2\mathbf{v}_2 \cos(2\varphi'), \quad \varphi' = \varphi - \varphi_R$

FOPI detector @ GSI



Systematics of Directed Flow & Stopping



Stopping:

$$b/b_{\max} < 0.15$$

$$\text{vartl} = \frac{\sigma^2(y_t)}{\sigma^2(y_z)}$$

Sideflow:


$$b/b_{\max} \simeq 0.3 - 0.4$$

$$\max [(p_x^{\text{dir}})^{(0)}]$$

W. Reisdorf et al., (FOPI), PRL 92 (2004) 232301

- Correlation between stopping & flow & pressure
- Evidence for incomplete stopping
- Stopping: maximum $\sim 400A$ MeV
 - decreasing towards higher beam energies
 - rising with system size, no saturation
 - below expectations from hydrodynamics

Systematics of Elliptic Flow



- **Transition** from in-plane to out-of-plane preferred emission at low energies
- **Maximum** $\sim 400A$ MeV (depending on Pt)
- v_2 **decreasing** toward higher beam energies


A. Andronic et al., (FOPI), GSI Report 2004-1 (2004) 54

- **Interplay between fireball expansion & spectator shadowing**
- **Passing time decreasing at high beam energies**
- **Influence of collision dynamics**
- **Information on different stages of the collision**

\Rightarrow High p_t particles messengers of high density phase

T. Gaitanos et al., Eur. Phys. J. A 12 (2001) 421


Shape parameters: Sensitivity to in-medium σ_{nn} ?



- $\theta_F \rightarrow$ Directed flow
- $\lambda_{31} = f_3^2/f_1^2 \rightarrow$ Directed flow & Stopping
- $\lambda_{21} = f_2^2/f_1^2 \rightarrow$ Elliptic flow

J. Gosset et al., (DIOGENE), Phys. Lett. B 247 (1990) 233

Ru (400 AMeV) + Ru - Proton-likes - $\langle b_{geo} \rangle = 1.1$ fm



N. Bastid et al., (FOPI), Nucl. Phys. A (2004), in press


Data favour in-medium σ_{nn} close or slightly higher than σ_{nn}^{free}

\Rightarrow Consistent with results on nuclear stopping

F. Rami et al., (FOPI), Phys. Rev. Lett. 84 (2000) 1120


B. Hong et al., (FOPI), Phys. Rev C 66 (2002) 034901

EoS from Directed Flow?



- Sensitivity to the EoS parametrization
- Soft EoS (with MDI & $\sigma_{nn}^{\text{free}}$) in best agreement with directed flow data for Au + Au & Xe + CsI at 400 A MeV
- Difficulties of the model to reproduce directed flow *versus* system size & low E_{beam} (90A MeV)

EoS from Elliptic Flow?



T. Kress, (FOPI), PhD, Darmstadt (2002)

- Proton elliptic flow in qualitative agreement with IQMD
 - Light fragments & IMF ($Z > 2$) abundantly produced at SIS energies
 - Bound protons/all protons:
 - ↪ 67% (400 A MeV) → 33% (1500 A MeV)
- ⇒ Total baryon elliptic flow not described by any EoS

Flow from Lee-Yang Zeroes method

Genuine flow directly from correlation between many particles

⇒ Non-flow correlations due to quantum statistics, resonance decays, momentum conservation effects, ..., not neglected

□ Generating function:

$$G(\mathbf{ir}) = \langle \prod_j [1 + \mathbf{ir} \omega_j \cos(n(\varphi_j - \theta))] \rangle_{\text{events}}$$

where $\ln G(\mathbf{ir}) = \sum_{k=1}^{+\infty} c_k \frac{(\mathbf{ir})^k}{k!}$, $c_k = \text{cumulant}$

□ Find first zeroe (minimum), r_0^θ , of $|G(\mathbf{ir})|$

$r_0^\theta \rightarrow$ Asymptotic behaviour of c_k in the expansion of $\ln G(\mathbf{ir})$

□ “Integrated” flow: $V_n^\theta\{\infty\} = \frac{j_{01}}{r_0^\theta}$ (& averaged over θ)

□ Resolution parameter: $\chi = \frac{V_n\{\infty\}}{\sigma}$

→ $\chi > 1$: Lee-Yang zeroes should be used

→ $0.5 < \chi < 1$: Important to optimize weights

→ $\chi < 0.5$: Large statistical errors, better to use cumulants

□ Differential flow:


→ Deduced from $V_n^\theta\{\infty\}$ in harmonics multiples of n

Detailed description in:


*R.S. Bhalerao et al., Phys. Lett. B 580 (2004) 157 & Nucl. Phys. A 727 (2003) 373;
N. Borghini et al., nucl-th/0402053 (2004)*

First application of Lee-Yang theory to FOPI data: Ru + Ru @ 1.69A GeV

Generating function



- $\chi = 1.45 \Rightarrow$ Lee-Yang Zeroes theory can be used
- Clear indication of collective effects



- Non-flow effects from 4-particle correlations negligible
- Evidence for (small) momentum conservation effects on v_1
- Non-flow effects negligible for higher harmonics

Ongoing development $\rightarrow \pi^\pm$ flow & influence of Δ decay?

(110 Millions central Ni + Ni @ 1.93A GeV)

Conclusion

Complete set of data at SIS energies measured with FOPI:

- Variation of **beam energy** from 90A MeV to 2A GeV
 - Variation of **system size** from Ca to Au
 - Variation of **asymmetry in isospin** (Ru/Zr)
 - Variation of **asymmetry in system size** (Au/Ca & Pb/Ni)
-
- Main dependences of directed & elliptic flow are available
 - New procedure of Lee-Yang Zeroes (& cumulants at SIS) successfully used for first time to analyze flow
 - Correlations from non-flow effects negligible for protons & composite particles
 - Most features of flow data reproduced qualitatively well by IQMD model but not in detail
-
- EoS is influencing different observables
 - EoS is linked to in-medium NN interaction
 - ⇒ momentum dependence, cross sections
 - Non-equilibrium effects important

FOPI Collaboration

A. Andronic, V. Barret, Z. Basrak, N. Bastid,
M.L. Benabderrahmane, R. Čaplar, E. Cordier, P. Crochet, P. Dupieux,
M. Dželalija, Z. Fodor, I. Gasparić, Y. Grishkin, O. Hartmann, N. Herrmann,
K.D. Hildenbrand, B. Hong, D. Kang, J. Kecskemeti, Y.J. Kim,
M. Kirejczyk, P. Koczon, M. Korolija, R. Kotte, M. Kowalczyk, T. Kress,
A. Lebedev, Y. Leifels, X. Lopez, A. Mangiarotti, V. Manko, T. Matulewicz,
M. Merschmeyer, D. Moisa, D. Pelte, M. Petrovici, F. Rami, W. Reisdorf,
A. Schuettauf, Z. Seres, B. Sikora, K.S. Sim, V. Simion,
K. Siwek-Wilczyńska, M. Smolarkiewicz, V. Smolyankin,
J. Soliwoda-Poddany, M. Stockmeier, G. Stoicea, Z. Tyminski,
K. Wiśniewski, D. Wohlfarth, Z. Xiao, I. Yushmanov, A. Zhilin

NIPNE Bucharest, Romania
KFKI Budapest, Hungary
LPC Clermont-Ferrand, France
GSI Darmstadt, Germany
Univ. of Heidelberg, Germany
IKH Rossendorf/Dresden, Germany
ITEP Moscow, Russia
Kurchatov Institute, Moscow, Russia
Korea University Seoul, Korea
IReS Strasbourg, France
RBI Zagreb, Croatia
Univ. of Warsaw, Poland

FOPI